



Volume 29, Issue 2, December 2015

Geospatial Assessment of Coseismic Landslides in Baturagung Area

Aditya Saputra, Munun Sartohadi, Danang Sri Hadmoko, Christopher Gomez

Benefits of Rural Biogas Implementation to Economy and Environment: Boyolali Case Study

Zetaria Tazi Hnyine, Saut Sagala, Wahyu Lubis, Dodon Yamin

Geothermal Energy Utilization in The Kamojang Nature Reserve, West Java, Indonesia (Pemanfaatan Energi Panas Bumi di Cagar Alam Kamojang, Jawa Barat, Indonesia)

Endang Nurfi Imarasa Dachlan, Vella Putiksari, Lilik Budi Prasetyo

Spatial Pattern of Bathymetry in Wangi-Wangi Island and its Surrounding Based on GEBCO Data and Nautical Map

Yulius, H L Salim, M Pamdhan

Geospatial Analysis and Turbidity Measurement for Monitoring Suspended Solids of Hitotsuse Dam in Miyazaki Prefecture, Kyushu, Japan

Purnama Budi Santosa, Yasuhiro Mitani

Development and Evaluation of Web GIS Application for Groundwater Management of Karanganyar Regency

Jumadi and Yuli Priyana

Morphometry Characteristics of Riverbed Sediment Grains as Basic Indicator Management of River Valley Environment (Case Study of Bogowonto River, Central Java)

Suorapto Dibyosaputro

Estimate of Changes in Carbon Stocks Based on Land Cover Changes in the Leuser Ecosystem Area (LEA) Indonesia

Ded Hermon

Accredited by the Minister of Research, Technology and Higher Education,
Republic of Indonesia, Based on the Decree Number 12/M/Kp/III/2015, Dated 11 February 2015

ISSN: 0852-0882 E-ISSN: 2460-3945



Accredited by DGHE, No. 12/14/10p/11/2015
Dated 11 Feb. 2015

Forum Geografi

Indonesian Journal of Spatial and Regional Analysis



Under Collaboration
between Fakultas Geografi UMS and
Ikatan Geograf Indonesia (IGI)
Published by MUP Press



- [Home](#)
- [About](#)
- [Login](#)
- [Register](#)
- [Search](#)
- [Current](#)
- [Archives](#)
- [Announcements](#)
- [In Press](#)
- [Ethics](#)


User Vol 29, No 2 (2015)

Language Forum Geografi

Journal Content Table of Contents

Notifications Articles

<p>Information ▼</p> <p>Keywords ▼</p> <p>Current Issue ▼</p>	<p>Geospatial Assessment of Cosmogenic Landslides in Batanguna Area Aditya Saputra, Jimun Sartohadi, Danang Sri Hadmoko, Christopher Gomez PDF 99 - 114</p> <p>Benefits of Rural Logistics Implementation to Economy and Environment: Boreilali Case Study Zakaria Tazi Hnyina, Saut Sagala, Wahyu Lubis, Dodor Yamin PDF 115 - 128</p> <p>Geotherm / Energy Utilization in the Kamojang Nature Reserve, West Java, Indonesia Endes N Dahlan, Mella Putiksari, Liik Budi Prasetyo PDF 129 - 143</p> <p>Spatial Pattern of Settlements in Wangi-Wangi Island and Its Surrounding, based on GEBCO (Global Nautical Map) Yulius Yulius, H L Salim, M Ramdhan PDF 144 - 152</p> <p>Geospatial Analysis and Turbidity Measurement for Monitoring Suspended Solids of Hirayasu (Tama) Miyazaki Prefecture, Kyushu, Japan Purnama Budi Samosa, Yasuhiro Mitani PDF 153 - 164</p> <p>Development and Evaluation of Web GIS Application for Groundwater Management of Karanganyar Regency Jumadi Jumadi, Yuli Priyana PDF 165 - 174</p> <p>Morphometry Characteristics of Fluvial Sediment Channels as Basic Indicator of Management of River Valley Environment (Case Study of Pagar Wonto River, Central Java) Suprpto Dibyosanturo PDF 175 - 187</p> <p>Estimate of Change in Carbon Stocks Based on Land Cover Change in the Lebak Ecosystem Area (LEA) Indonesia Dedi Hermon PDF 188 - 196</p>
--	--



Volume 29, Issue 2, December 2015

ISSN 0852-0682



E-ISSN 2460-3945

Editor in Chief

Jumadi, S. Si., M. Sc.

Co-Editor in Chiefs

Drs. Yuli Priyana, M. Si.

Agus Anggoro Sigit, S. Si, M. Sc.

Drs. Priyono, M. Si.

Editorial Board Members

Dr. Kuswaji Dwi Priyono, M. Si. (*Geomorphology*) - UMS

Dr. Imari Harjono, M. Si. (*Environmental Analysis*) - UMS

Dra. ALif Nur Anna, M. Si. (*Water Resources*) - UMS

Drs. M. Musyam, M. TP. (*Human Geography*) - UMS

Dr. Sutarno, M. Si. (*Socio-economic*) - UMP

Dr. Sigit Heru Murti (*Remote Sensing Application*) - UGM

Aditya Saputra, S. Si, M. Sc. (*GIS Application*) - UMS

Amin Sunardi, S. Si, M. P. (*Spatial Analysis*) - UMS

Advisory Board Members

Prof. Dr. Suratman Woro Suprojo, M. Sc. (Universitas Gadjah Mada)

Dr. Christopher Gomez. (University of Canterbury, NZ)

Prof. Dr. Sudarmadji, M. Eng. (Universitas Gadjah Mada)

Prof. Dr. Rianta, M. Sc. (Universitas Gadjah Mada)

Prof. Dr. Junun Hartohadi, M. Sc. (Universitas Gadjah Mada)

Prof. Dr. Hartono, DEA, DESS. (Universitas Gadjah Mada)

Prof. Dr. Saibyalto, MS. (Universitas Gadjah Mada)

Prof. Dr. M. Agus Marfal, M. Sc. (Universitas Gadjah Mada)

Dr. Djati Marliyatro, M. Sc. (Universitas Gadjah Mada)

Prof. Dr. Sunarto, M. Si. (Universitas Gadjah Mada)

Prof. Dr. Chatusa Muryani, M. Si. (Universitas Sebelas Maret)

Dr. Djoko Hurmantyo, M. Si. (Universitas Indonesia)

Dr. Saat Saatla, M. Sc. (Institut Teknologi Bandung)

Publishing Manager

Choirul Amir, S. Si., M. M.

Publishing Period

Issued Period: Juli dan Desember

First Issue: Juli 1987

Publishing Institution



Geography Faculty
UNIVERSITAS MUHAMMADIYAH
SURAKARTA



IGI
INDONESIAN GEOGRAPHICAL
SOCIETY

Mailing Address:

Fakultas Geografi Universitas Muhammadiyah Surakarta

Jl. A. Yani Pabelan Kartasura Terminal Pos I Surakarta 57102, Telp (0271) 717417 Psw 151-153

Fax: (0271) 715448, website: journals.ums.ac.id or publikasiilmiah.ums.ac.id,

e-mail: forgeo@ums.ac.id

Contents

- Geospatial Assessment of Coseismic Landslides in Baturagung Area**
Aditya Saputra, Junun Sartohadi, Dianang Sri Hadmoko, Christopher Gomez 99-113
- Benefits of Rural Biogas Implementation to Economy and Environment: Boyolali Case Study**
Zakaria Tazi Hnyine, Saut Sagala, Wahyu Lubis, Dodon Yamin 114-127
- Geothermal Energy Utilization in the Kamojang Nature Reserve, West Java, Indonesia (Pemanfaatan Energi Panas Bumi di Cagar Alam Kamojang, Jawa Barat, Indonesia)**
Endes Nurfilmarasa Dachlan, Vello Putiksari, Lilik Budi Prasetyo 128-142
- Spatial Pattern of Bathymetry in Wangi-Wangi Island and its Surrounding Based on GEBCO Data and Nautical Map**
Yulius, H L Salim, M Ramdhan 143-151
- Geospatial Analysis and Turbidity Measurement for Monitoring Suspended Solids of Hitotsuse Dam in Miyazaki Prefecture, Kyushu, Japan**
Purnama Budi Santosa, Yasuhiro Mtani 152-163
- Development and Evaluation of Web GIS Application for Groundwater Management of Karanganyar Regency**
Jumadi and Yuli Priyana 164-173
- Morphometry Characteristics of Riverbed Sediment Grains as Basic Indicator Management of River Valley Environment (Case Study of Bogowonto River, Central Java)**
Suprpto Dibyosaputro 174-186
- Estimate of Changes in Carbon Stocks Based on Land Cover Changes in the Leuser Ecosystem Area (LEA) Indonesia**
Dedi Hermon 187-196

Estimate of Changes in Carbon Stocks Based on Land Cover Changes in the Leuser Ecosystem Area (LEA) Indonesia

Dedi Hermon

Geography Department

and Graduate Program State University of Padang Indonesia

Corresponding E-mail: dihermon006@gmail.com

Abstract

This research aimed at designing the model of land cover changes in 1990 and 2014, and estimating carbon stock changes in each land cover in Leuser Ecosystem Area (LEA). The spatial model of land cover changes was analyzed by interpreting Landsat 5 TM imagery in 1990 and Landsat 7 ETM+ imagery in 2014 with ERDAS 9.1 and Land Change Modeller (LCM) in Idrisi TerrSet v.18. The analysis of land area changes (ha) in each land cover from 1990 to 2014 used ERDAS 9.1 with tools Interpreter (GIS Analysis-Matrix). Systematic survey method was employed in order to analyze carbon stocks. The sampling technique was stratified purposive composite sampling which used plot technique. The estimate of tree biomass used allometric equation. The estimate of carbon stocks in each land cover in 1990 was measured based on the total of carbon stocks in 2014 which was converted with the areas of each land cover in 1990. Spatial model of land cover changes in LEA in 1990-2014 showed the changes of area in each land cover which caused the changes of carbon stocks in each land cover as well.

Keywords: Leuser Ecosystem Area, Carbon Stocks, Land Cover Change

Abstrak

Tujuan penelitian adalah untuk menyusun model perubahan tutupan lahan tahun 1990 dan 2014 serta melakukan estimasi perubahan cadangan karbon pada setiap tutupan lahan di Leuser Ecosystem Area (LEA). Model spasial perubahan tutupan lahan LEA di analisis dengan melakukan interpretasi Citra Landsat 5 TM tahun 1990 dan Citra Landsat 7 ETM+ tahun 2014 dengan ERDAS 9.1 dan Land Change Modeller (LCM) dalam Idrisi TerrSet v.18. Analisis perubahan luas lahan (ha) pada masing-masing tutupan lahan periode tahun 1990-2014 menggunakan ERDAS 9.1 dengan tools Interpreter (GIS Analysis-Matrix). Analisis cadangan karbon menggunakan metode survey secara sistematis dengan teknik pengambilan sampel adalah stratified purposive komposit sampling dengan menggunakan teknik plot. Adapun pendugaan biomassa pohon menggunakan persamaan allometrik. Penghitungan perkiraan cadangan karbon pada masing-masing tutupan lahan pada tahun 1990 adalah berdasarkan hasil perhitungan karbon pada tahun 2014 yang dikonversikan dengan luasan masing-masing tutupan lahan pada tahun 1990. Model spasial perubahan tutupan lahan di Leuser Ecosystem Area (LEA) periode 1990-2014, menunjukkan terjadinya perubahan luas (ha) pada masing-masing tutupan lahan, sehingga mengakibatkan terjadinya perubahan cadangan karbon pada masing-masing tutupan lahan.

Keywords: Leuser Ecosystem Area, Carbon Stocks, Land Cover Change

Introduction

Carbon is closely related to land cover changes because the largest carbon stocks are on forest area. Hermon (2012a) states that carbon stocks in forest land cover are relatively bigger than any other land covers namely shrub, cultivation, and paddy field. Loss of carbon stock happens if land cover

of forest, cultivation, and paddy field are changed into settlements which make the dynamics of carbon stock in each land cover are decreasing year by year. Harun (1992); Kustiawan (1997); Pribadi *et al.* (2006); Hermon (2009); and Hermon (2012a) explain that the changes of land cover are mostly triggered by people economic activities and development which then dominate and impose

natural areas (forest) to change their function. This condition makes the decreasing of carbon stocks in certain areas which then reduce more carbon stocks every year.

Lusiana *et al.* (2005); Hermon (2010); Hermon (2012a); and Hermon (2012b) elaborate that carbon has much influence on climate change, and can give negative effect to the earth because 20% of greenhouse gasses are caused by deforestation or the change of forest areas into another type of land cover. In 2000, it was estimated that the land use emissions, and the changes of land use in Indonesia was 2, 563 Mt CO₂ or equal to 20% of the total emissions of land and forest changes in the world. These emissions were mostly caused by deforestation and forest degradation. Hermon (2014a) explains about the correlation between land cover changes and carbon stocks, and its impact on climate change in Padang. It was explained that the land cover changes of forest, shrubs, cultivation to settlements in Padang from 1985-2011 triggered the increasing of greenhouse gases concentrate especially the increasing amount of CH₄, CO₂, and N₂O. The increasing amount of CO₂ was caused by forest logging and burning for becoming settlements or other economic activities.

The changes of forest land cover into other types of land cover also happened in Leuser Ecosystem Area (LEA). From 2006 to 2009, there was a decrease of LEA forest area for ± 400,00 ha in Southeast Aceh Regency. This decrease indicated that LEA was having damaged because of the conversion of forest land cover to other types of land cover. The changes that occurred from year to year cause the change of carbon stocks (WWF, 2015). LEA itself is rainforest areas and national parks which consist of swamps, lowland and upland forests, and mountains which are located in two provinces-Aceh and North Sumatra. Geographically, LEA lies between Laut Tawar Lake in Aceh Province and Toba Lake in North Sumatra Province with 2.6 million hectares areas which cover Southeast Aceh Regency, Gayo Lues, Central Aceh, Bener Meriah, Aceh Singkil, Kota Subulussalam,

South Aceh, Southwest Aceh, Nagan Raya, West Aceh, Aceh Tamiang, East Eceh, North Aceh, Langkat, Dairi, Karo and Deli Serdang. According to its geomorphological areas, LEA has slopes from flat to very steep ones, with the rainfall between 2500-4000 mm/year. LEA has the biggest biodiversity in Asia, so that it has many ecological functions such as water reserve, climate stabilization, mitigation of erosion and flash floods, prevention of pest outbreaks, and carbon stocks (Forestry Department and IFCA, 2007; WWF, 2015; Hermon 2014b; Hermon, 2014c).

LEA has significant role for keeping the stability of earth temperature and climate; thus, LEA functions as National Park, and is also determined as Biosphere reserves by UNESCO. Threat of LEA sustainability is the conversion of land to become agricultural land (cultivation and paddy field), plantation, and settlements for people living around LEA. This land conversion will change forest land cover into agricultural and residential land cover in which this condition will directly change the carbon stocks in LEA (TFCA Sumatera, 2015). The change of carbon stocks are analyzed based on the change of land cover by combining analysis technique of remote sensing imagery and field survey to measure above ground biomass. The objective is to formulate the spatial model of LEA land cover changes as guidance in estimating carbon stock from 1990 to 2014.

Research Method

Spatial Model of Land Cover Changes in Leuser Ecosystem Area (LEA)

The model of land cover change is analyzed based on Zain (2002); Hermon (2009); and Hermon (2012a) by interpreting Landsat 5 TM 1990 imagery and Citra Landsat 7 ETM+ 2014 (path row 130-58) with ERDAS 9.1 as the analysis tool. The classification of land cover change is analyzed with supervised classification technique for each image which resulted on six temporary patterns of land cover, namely (1) forest, (2) cultivation, (3)

paddy field, (4) shrub, (5) plantation, (6) settlement, and (7) body of water. Before classifying land cover by using supervised classification technique, separability analysis in the form of quantitative analysis is carried out in order to inform us the evaluation of separated areas in each class as a validation of band combination using ENVI 4.5 (Hermon, 2009). The analysis of land cover change is conducted by using Land Change Modeller (LCM) Idrisi TerrSet v.18. Meanwhile, the analysis of land area change (hectares) in each land cover in 1990 and 2014 is conducted by using ERDAS 9.1 with tools Interpreter (GIS Analysis-Matrix).

The Analysis of Carbon Stock Change in Leuser Ecosystem Area (LEA)

The analysis of carbon stock in each land cover is carried out by using Above Ground Biomass (AGB) technique with systemic survey method. Sampling technique used is stratified purposive composite sampling with plot technique (Hairiah and rahayu, 2007; Hermon, 2012a). The plot size for taking the sample of trees is 10x10m; meanwhile 1x1 m of plot size which is made in each land cover is used for taking the sample of undergrowth and litter (Hermon, 2012a). The samples of trees (stalks, branches, twigs, and leaves) are taken compositely and analyzed with non-destructive method; meanwhile the samples of undergrowth and litter are analyzed with destructive method. The estimate of tree biomass uses allometric equation (Kattering, 2001) with $BK = 0.11 p D^{2.62}$ in which D is the diameter of tree at breast height, cm) and p is the density of wood. The biomass

of undergrowth and litter is measured based on the total dry weight per quadrant (in gr/m² and conversed to ton/ha) which uses the formula: Total of BK (gr) = BK/BB x total of BB (gr), in which BK (dry weight, gr) and BB (wet weight, gr). The total of carbon stocks is got from the total of biomass multiplied with 0.46 (the concentrate of C in organic materials is about 46%) (Hairiah and Rahayu, 2007; Hermon, 2012a). The estimate of carbon stocks in each land cover in 1990 was based on the estimate of carbon stocks in 2014 which was conversed with the areas of each land cover in 1990 (Hermon, 2012a).

Result and Discussion

Spatial Model of Land Cover Change in Leuser Ecosystem Area (LEA)

The analysis using Landat 5 TM imagery in 1990 and Citra Landsat 7 ETM+ imagery in 2014 resulted on the data of land cover changes for the last 24 years, between 1990 and 2014. Spatial model of land cover change in LEA from 1990 to 2014 can be seen in Picture 1 below. The spatial model of land cover change in LEA from 1990 to 2014 informed us that land cover for settlements was increasing in 2014 compared to 1990; on the other hand, land cover of forest in 2014 as well as land cover of cultivation, plantation, and shrub was relatively decreasing compared to 1990. The increasing of land cover area for paddy field occurred in 2014 compared to 1990. Baban and Yusof (2001); Amdam (2004); and Antrop (2004) explain that spatial model of land cover change will give the information spatially about all variation of land cover changes.




Figure 1. Spatial Model of Land Cover Change in The Leuser Ecosystem Area (LEA) 1990 and 2014

The variation of land cover changes in LEA in 1990 and 2014 was caused by the activities of people surrounding LEA, such as for enhancing settlements, for new paddy field areas, or other economic activities (TFCA, 2015). The reduction of forest area and the

increasing of settlements in LEA indicate that people activities are no longer controllable in utilizing land cover. The reduction of forest area can be seen in the area between South Aceh Regency and Southwest Aceh (WWF, 2015).




Figure 2. Gains and Losses Land Cover Area in The Leuser Ecosystem Area (LEA) in Year Period 1990-2014

The data reveal that in 1990 the area of forest in LEA was 1,341,322.1 ha; cultivation was 920,528.3 ha; settlement was 153,231.1 ha; and the body of water was 1,576.1 ha. Compared to 2014, there was changes in each land cover in which the area of forest was decreasing into 1,205,576.2 ha; cultivation was decreasing into 638,331.1 ha; paddy field was increasing into 247,319.1 ha; shrub was decreasing into 99,098.1 ha; plantation was decreasing into 19,333.2 ha; and settlement was increasing into 388,766.3 ha; meanwhile, there was no change on the body of water which was 1.576.1 ha. Chust et al (2004) conveys that

no changes in the body of water in certain period of time indicate that the changes of land cover are mostly oriented on agriculture and new settlements. Furthermore, Hermon (2012a) elaborates that land cover changes into agriculture (paddy field, cultivation) and settlements have big influence on the changes and loss of carbon stocks. The decreasing of carbon stocks happen if there is a conversion of land cover of forest to agricultural area, and the loss of carbon stocks happen when land cover of forest, paddy field, and shrub is converted into settlements

Table 1. Land Cover Change in the Leuser Ecosystem Area (LEA) in Year Period 1990-2014

No	Land Cover	1990 (ha)	2014 (ha)	Extensive Changes in Land Cover (ha)	
				Increase	Reduce
1	Forest	1.341.322,1	1.205.576,1	0	135.746,0
2	Cultivation	920.528,3	638.331,1	0	282.197,2
3	Paddy Field	19.113,0	247.319,1	228.206,1	0
4	Shrub	128.575,3	99.098,1	0	29.477,2
5	Plantation	35.654,1	19.333,2	0	16.320,9
6	Settlement	153.231,1	388.766,3	235.535,2	0
7	Body of Water	1.576,1	1.576,1	0	0
	Total	2.600.000	2.600.000	463.741,3	463.741,3

Source: Result of GIS Matrix Analysis of Landsat 5 TM 1990 and Landsat 7 ETM+ 2014 using ERDAS 9.1 and LCM Idrisi TerrSet v.18 (2015)

In the period of 1990-2014, there were many changes of land cover areas into settlements such as forest areas which became settlements for 98,254.3 ha, cultivation area becoming settlements for 132,380.1 ha, paddy field area

becoming settlements for 876.1 ha, shrub area becoming settlements for 578.5 ha, and the changes of plantation areas becoming residential areas for 3,445.2 ha.




Figure 3. Land Cover Changes into Settlements in The Ecosystem Area (LEA) Period 1990-2014

Fitzsimmons (2003); Lubowski *et al.* (2006); Ritohardoyo (2007); Hermon (2009); and Hermon (2012a) explain that land cover changes especially the changes of land cover of forest into settlements will continue to occur and always spatially varied either planned or unplanned. Dahroni (2008) states that settlement will always be a source of problem in human history since the development of settlements will cause the uncontrollable land cover changes, particularly on the changes

of land cover of forest into other land covers related to settlements and economic activities.

The Estimate of Carbon Stock Changes in Each Land Cover \in Leuser Ecosystem Area (LEA)

The changes of forest, cultivation, paddy field, shrub, and plantation areas into settlements in LEA from 1990 to 2014 indicated the change of carbon stocks in LEA. For more detail, it is shown in Table 2 below.

Table 2. Biomass and Carbon Stocks in the Leuser Ecosystem Area (LEA) in 2014

Land Cover	Trees Number/ha	Tree Biomass		Litter Biomass		Total Biomass (ton/ha)	Carbon (ton/ha)
		(kg/m ²)	(ton/ha)	(kg/m ²)	(ton/ha)		
Forest	341	365,22	3.652,20	18,320	183,20	3.835,40	1.764,28
Cultivation	57	15,77	157,70	15,44	154,40	312,10	146,56
Paddy Field	0	0,00	0,00	19,89	198,90	198,90	91,26
Shrub	16	2,87	28,70	26,93	269,30	298,00	137,08
Plantation	239	64,84	648,40	7,11	71,10	719,50	330,97
Settlement*	--	--	--	--	--	--	--
Body of Water*	--	--	--	--	--	--	--
Total	653	448,70	4487,00	87,69	876,90	5.363,90	2.470,15

Source: Data Analysis Research, 2015

Based on the condition in 2014, the carbon stocks per hectare in LEA was 2,470.15 ton/ha which spread on forest area for 1,764.28 ton/ha, paddy field for 91.26 ton/ha, shrub for

137.08 ton/ha, and plantation for 330.97 ton/ha. The total of carbon stocks in 2014 in LEA is presented in Table 3 below.

Table 3. Total Carbon Stocks in The Leuser Ecosystem Area (LEA) 2014

Land Cover	Area (ha)	Carbon Stocks (ton/ha)	Total Carbon Stocks (ton)
Forest	1.205.576,1	1.764,28	2.126.973.801,70
Cultivation	638.331,1	146,56	93.553.806,01
Paddy Field	247.319,1	91,26	22.570.359,31
Shrub	99.098,1	137,08	13.584.367,54
Plantation	19.333,2	330,97	6.398.874,68
Settlement*	388.766,3	--	--
Body of Water*	1.576,1	--	--
Total	2.600.000	2.470,15	2.263.081.209,24

Source: Data Analysis Research, 2015

* Not Analyzed

In 2014, the total of carbon stocks in LEA was 2,263,081,209.24 ton which spread on forest area for 2,126,973,801.70 ton, cultivation for 93,553,806.01 ton, paddy field for 22,570,359.31 ton, shrub for 13,584,367.54

ton, and plantation 6,398,874.68 ton. The biggest carbon stocks were in the forest so that the preservation of forest area must be sustainable. The conversion result of carbon stocks in LEA in 1990 can be seen in Table 4.

Table 4. Prediction of Total Carbon Stocks in the Leuser Ecosystem Area (LEA) 1990

Land Cover	Area (ha)	Carbon Stocks Prediction (ton/ha)	Prediction of Total carbon Stocks (ton)
Forest	1.341.322,1	1.764,28	2.366.468.283,87
Cultivation	920.528,3	146,56	134.912.686,27
Paddy Field	19.113,0	91,26	1.744.252,38
Shrub	128.575,3	137,08	17.625.102,13
Plantation	35.654,1	330,97	11.800.437,48
Settlement*	153.231,1	--	--
Body of Water*	1.576,1	--	--
Total	2.600.000	2.470,15	2.532.550.762,13

Source: Data Analysis Research, 2015

* *Not Analyzed*

Based on the conversion result of carbon stocks, it was known that the carbon stocks in LEA in 1990 were 2,532,550,762.13 ton. In details, forest areas had the carbon stocks for 2,366,468,283.87 ton, cultivation for 134,912,686.27 ton, paddy field for

1,744,252.38 ton, shrub for 17,625,102.13 ton, and plantation for 11,800,437.48 ton. In the period of 1990-2014, there was decrease and increase of carbon stocks on each land cover, as shown in Table 5.

Table 5. Decrease and Increase Carbon Stocks on Each Land Cover in the Leuser Ecosystem Area (LEA) Period 1990-2014

Land Cover	Area (ha) 1990	Prediction of Total Carbon Stocks (ton/ha)	Area (ha) 2014	Total Carbon Stocks (ton)	Decrease in Carbon Stocks (ton)	Increase Carbon Stocks (ton)
Forest	1.341.322,1	2.366.468.283,87	1.205.576,1	2.126.973.801,70	239.494.482,17	0,0
Cultivation	920.528,3	134.912.686,27	638.331,1	93.553.806,01	41.358.880,26	0,0
Paddy Field	19.113,0	1.744.252,38	247.319,1	22.570.359,31	0,0	20.826.106,93
Shrub	128.575,3	17.625.102,13	99.098,1	13.584.367,54	4.040.734,59	0,0
Plantation	35.654,1	11.800.437,48	19.333,2	6.398.874,68	5.401.562,80	0,0
Settlement*	153.231,1	--	388.766,3	--	--	--
Body of Water*	1.576,1	--	1.576,1	--	--	--
Total	2.600.000	2.532.550.762,13	2.600.000	2.263.081.209,24	290.295.659,82	20.826.106,93

Source: Data Analysis Research, 2015

* *Not Analyzed*

The decrease of carbon stocks in LEA in 1990-2014 occurred in forest land cover (239,494,482.17 ton), cultivation (41,358,880.26), shrub (4,040,734.59 ton), and plantation (5,401,562.80 ton). On the contrary, the increase of carbon stocks occurred in land cover of paddy field (20,826,106.93 ton) because land cover of paddy field was

increasing in 2014 for 228,206.1 ha. The loss of carbon stocks in LEA because of the changes of land cover into settlements from 1990 to 2014 was quite big; it was 194,049,235.370 ton. In details, the loss of carbon stocks because of land cover changes into settlements can be seen in Table 6.


Table 6. Loss of Carbon Stocks in Leuser Ecosystem Area (LEA) because of Land Cover Changes into Settlements

No	Land Cover Change	Area (ha)	Losses of Carbon Stocks (ton)
1	Forest-Settlement	98.254,3	173.348.096,404
2	Cultivation-Settlement	132.380,1	19.401.627,456
3	Paddy Field-Settlement	876,1	79.952,886
4	Shrub-Settlement	578,5	79.300,780
5	Plantation-Settlement	3.445,2	1.140.257,844
	Total	235.534,2	194.049.235,370

Source: Result of GIS Matrix Analysis of Landsat 5 TM 1990 and Landsat 7 ETM 2014 using ERDAS 9.1 and LCM Idrisi TerrSet v.18 (2015) and Data Analysis Research (2015)

In the period of 1990-2014, there were many changes of land cover areas into settlements in LEA such as forest areas which became settlements for 98,254.3 ha. It means that the loss of carbon stocks in forest area was 173,348,096,404 ton. The cultivation area becoming settlements was for 132,380.1 ha, which indicated 19,401,627,456 ton of

carbon stock loss. The land cover changes also occurred in paddy field area in which it became settlements for 876.1 ha which means 79,952,886 ton carbon stock loss. Then, the changes of plantation areas becoming residential areas were for 3,445.2 ha, and it indicated the loss of carbon stocks for 1,140,257.844 ton.

**Figure 4. Carbon Loss in The Leuser Ecosystem Area (LEA)**

It was found that the biggest carbon stock loss in LEA happened because of the land cover changes of forest areas into settlements for 173,348,096.404 ton from 1990 to 2014. Hermon (2012a) explains that the loss of carbon stocks was triggered by land cover changes into settlements which kept increasing through conversion of forest, shrub, and paddy field

into settlements. Hermon (2014a) elaborates that these land cover changes into settlements cause the removal of carbon stocks in the forest into the atmosphere which then become CO₂ through deforestation and land burning. This condition will cause the accumulation of CO₂ in the atmosphere.

Conclusion

Land cover changes occurred because of people activities which are related to economy and settlements. The changes of land cover will directly change the amount of carbon stock in an area. From 1990-2014, there was a big change of land cover happened in LEA. Land cover of paddy field and settlements were increasing a lot in 2014; meanwhile, land cover of forest, cultivation, shrub, and

plantation was decreasing in 2014. In addition, the changes of land cover becoming settlement were the changes dominantly happened in LEA compared to other land cover. This condition caused the decrease of carbon stocks in LEA in 2014 compared to carbon stocks in 1990 for each land cover. The big changes of either land cover of forest or other land covers into settlements had cause the loss of carbon stocks in LEA for 194,049,235.370 ton from 1990 to 2014.

References

- Antrop, M. (2004). Landscape Change and the Urbanization Process in Europe. *Landscape and Urban Planning*. 67: 9–26. www.elsevier.com/landurbplan
- Amdam, R. (2004). Spatial County Planning as a Regional Legitimizing Process. *European Journal of Spatial Development*. 11: 1-22 <http://www.nordregio.sc/EJSD>
- Baban, S.M., and K.W. Yusof. (2001). Mapping Land Use/Cover Distribution on a Mountainous Tropical Island Using Remote Sensing and GIS. *International Journal of Remote Sensing*. 22: 1909-1918
- Chust, G., D. Ducrot, and J.L.I. Pretus. (2004). Land Cover Mapping with Patch-Derived Landscape Indices. *Landscape and Urban Planning* 67: 45-53. www.elsevier.com/locate/landurbplan
- Dahroni. (2008). Analisis Keruangan terhadap Perubahan dan Persebaran Permukiman Kumuh di Kecamatan Banjarsari Kota Surakarta Tahun 2001-2005. *Forum Geografi*. Vol. 22. No. 1. Juli 2008: 85-96
- Fitzsimmons, M. (2003). Effects of Deforestation and Reforestation on Landscape Spatial Structure in Boreal Saskatchewan, Canada. *Forest Ecology and Management*. 174: 577-592. www.elsevier.com/locate/foreco
- Departemen Kehutanan dan IFCA. (2007). REDDI, Reducing Emissions from Deforestation and Forest Degradation in Indonesia, REDD Methodology and Strategies Summary for Policy Makers. Dephut. Jakarta
- Hairiah, K dan Rahayu, S. (2007). Pengukuran “Karbon Tersimpan” di Berbagai Macam Penggunaan Lahan. Bogor. World Agroforestry Centre – ICRAF, SEA Regional Office, University of Brawijaya, Unibraw, Indonesia.
- Harun, U.R. (1992). Dinamika Penggunaan Sumberdaya Lahan di Jawa Barat 1970-1990. *Jurnal PWK*. 3: 48-53
- Hermon, D. (2009). Dinamika Permukiman dan Arah Kebijakan Pengembangan Permukiman pada Kawasan Rawan Longsor di Kota Padang. Disertasi. IPB Bogor
- Hermon, D. (2010). *Geografi Lingkungan: Perubahan Lingkungan Global*. UNP Press. Padang
- Hermon, D. (2012a). Dinamika Cadangan Karbon akibat Perubahan Tutupan Lahan menjadi Lahan Permukiman di Kota Padang Sumatera Barat. *Forum Geografi*. Vol. 26. No. 1, Juli 2012: 29-44
- Hermon, D. (2012b). Mitigasi Bencana Hidrometeorologi: Longsor, Banjir, Degradasi Lahan,

Ekologi, Kekeringan, dan Puting Beliung. UNP Press. Padang

- Hermon, D. (2014a). Impacts of Land Cover Change on Climate Trend in Padang Indonesia. *Indo J. Geog.* Vol. 46. No. 2: 104-209
- Hermon, D. (2014b). Arahan Mitigasi Bencana Longsor Kawasan Gunung Padang Kota Padang Sumatera Barat. *Jurnal Geografi.* Vol. 3. No. 2: 1-93
- Hermon, D. (2014c). *Geografi Bencana Alam.* Radja Grafindo Persada Press. Jakarta
- Ketterings, Q.M., Coe R., Meine van Noordwijk., Ambagau Y., Palm C.A. (2001). Reducing Uncertainty in the of Allometric Biomass Equation for Predicting Aboveground Tree Biomass in Mixed Secondary Forest. *Forest Ecology and Management* 146: 199-209
- Kustiawan, I. (1997). Permasalahan Konversi Lahan Pertanian dan Implikasinya terhadap Penataan Ruang Wilayah. Studi Kasus: Wilayah Pantura Jawa Barat. *Jurnal PWK.* 8: 49-60
- Lubowski, R.J., S. Bucholtz., R. Claassen., M.J. Roberts., J.C. Cooper., A. Gueorguieva., and R. Johansson. (2006). Environmental Effects of Agricultural Land-Use Change. The Role of Economics and Policy. A Report from the Economic Research Service. USDA. www.ers.usda.gov
- Lusiana, Betha., Meine van Noordwijk., Surbukti Rahayu. (2005). Cadangan Karbon di Kabupaten Nunukan Kalimantan Timur: Monitoring Secara Spasial dan Pemodelan. ICRAF. Bogor
- Pribadi, D.O., D. Shiddiq, dan M. Ermyanila. (2006). Model Perubahan Tutupan Lahan dan Faktor-Faktor yang Mempengaruhinya. *Jurnal Teknologi Lingkungan.* Pusat Pengkajian dan Penerapan Teknologi Lingkungan. 7: 35-51
- Ritohardoyo, S. (2007). Perubahan Permukiman Pedesaan Pesisir Kabupaten Gunung Kidul Daerah Istimewa Yogyakarta Tahun 1996-2003. *Forum Geografi.* Vol. 21. No. 1. Juli 2007: 78-92
- TFCA-Tropical Forest Coservation Action-Sumatera. (2015). Program Restorasi dan Konservasi Kawasan 13 Bentang Alam Prioritas di Sumatera. TFCA Sumatera. Jakarta
- WWF. (2015). Deforestation, Forest Degradation, Biodiversity Loss and CO₂ Emissions in Sumatra, Indonesia. WWF Indonesia Technical Report. Jakarta.
- Zain, A.F.M. (2002). Distribution, Structure dan Function of Urban Green Space in Southeast Asian Mega-Cities with Special Reference to Jakarta Metropolitan Region (JABOTABEK). Doctoral Degree Program. Department of Agricultural and Environmental Biology Graduate School of Agricultural and Life Sciences. The University of Tokyo. Japan